

Patient-mix Coefficients for October 2011 Publicly Reported HCAHPS Results

As noted in the HCAHPS *Quality Assurance Guidelines, V6.0*, prior to public reporting, hospitals' HCAHPS results are adjusted for the effects of both mode of survey administration and patient-mix. Generally speaking, HCAHPS adjustments for survey mode are larger than adjustments for patient-mix. The survey mode adjustments that are used in publicly reported HCAHPS results are reported in the paper entitled, "[Mode and Patient-mix Adjustment of the CAHPS[®] Hospital Survey \(HCAHPS\)](#) April 30, 2008," which can be found on www.hcahpsonline.org.

In order to derive the mode adjustment coefficients, it was necessary to conduct a randomized mode experiment. The resulting mode adjustment coefficients will not change as a function of the data used in public reporting. For more information on how the HCAHPS mode experiment was conducted and the survey mode and patient-mix adjustments were derived, please see, "*Effects of Survey Mode, Patient Mix, and Nonresponse on CAHPS Hospital Survey Scores.*" Elliott, M.N., A.M. Zaslavsky, E. Goldstein, W. Lehrman, K. Hambarsoomian, M.K. Beckett, and L. Giordano. *Health Services Research*. 2009. 44: 501-518.

The mode experiment data were also used to develop and validate the HCAHPS patient-mix model (which is referred to as "case-mix" elsewhere in the CAHPS literature), as described in the document referenced above. However, in the case of patient-mix adjustment, a randomized experiment is not necessary to accurately estimate the coefficients of the model.

In order to estimate the exact patient-mix coefficients as accurately as possible, we employ the large sample size of each quarterly national publicly reported data set. This approach allows us to detect changes in the association of patient-mix adjustors and HCAHPS measures over time and then adjust accordingly. This approach is consistent with recommended CAHPS practice for case-mix adjustment (www.cahps.ahrq.gov <<http://www.cahps.ahrq.gov/>>).

Patient-mix adjustment is performed within each quarter of data after data cleaning and before mode adjustment. *Coefficients* obtained in linear regression models (not reported) estimate the tendency of patients to respond more positively or negatively. The *adjustments* needed to counter that tendency are obtained by multiplying the patient-mix *coefficients* by (-1.0). Tables 1 and 2 below report patient-mix adjustments for the "top-box" (most positive response) and "bottom-box" (least positive response) respectively of the ten publicly reported HCAHPS measures (six composites, two individual items, and two global items), averaged across the four reported quarters.

As an example, patients aged 55-64 were 5.33% more likely to provide the most positive response (“Always”) for items in the *Communication with Nurses* composite when compared to the reference group of patients 85 and older. Thus, the corresponding adjustment for patients aged 55-64 relative to patients 85 and older for that composite is a subtraction of 5.33%, reflected in the “-5.33%” entry in Table 1. Similarly, for each level of decreasing self-rated health status (where 5=poor, 4=fair, 3=good, 2=very good, and 1=excellent), the percentage of patients providing an “Always” response for *Communication with Doctors* decreased by 5.00%. Thus, a patient in fair health (4) would have a $(4-1)*5.00\%=15.00\%$ lower chance of an “Always” response than a patient in excellent health (1), and the corresponding adjustment for a patient in poor health relative to a patient in excellent health would be +20.00%.

Publicly reported HCAHPS scores are adjusted to the overall national mean of patient-mix variables across all hospitals reporting in a given quarter (as reported in Table 3). Thus, whether the scores of a given hospital are adjusted upward or downward for a given measure depends not only on these patient-mix adjustments, but also on the patient-mix of that hospital relative to the national average of these patient-mix characteristics. Specifically, the total patient mix-adjustment for a given hospital is the sum of a series of products, where each product multiplies the adjustment in Table 1 (top-box) or Table 2 (bottom-box) by the deviation of the hospital’s mean on the corresponding patient-mix variable from the national mean on that patient-mix variable (from Table 3).

Four sets of numbers are needed to calculate final patient-mix adjusted scores for a given hospital: (1) Means of HCAHPS outcomes (top-box proportions or bottom-box proportions) for the hospital in question that have been adjusted for survey mode; (2) individual-level patient-mix adjustments from Tables 1 and 2 of this document; (3) that hospital’s means on patient-mix variables; and (4) national means on patient-mix variables from Table 3 of this document.

Below we provide additional detail regarding the calculation of the *response percentile* and *service line by age interaction* variables.

A hospital’s patient-mix adjustment variable *response percentile* is calculated as follows: For a given hospital and a given month, all completed surveys are ranked based on their respective “lag times.” Lag time is the number of days between a patient’s discharge from the hospital and the return of the mail survey, or the final disposition of the telephone or IVR survey. Ranks are averaged in the case of ties. Response percentile is calculated by dividing **lag time rank** by **monthly sample size**.

The *service line by age interaction* variables used in patient-mix adjustment can be calculated by following the steps below for all completed surveys:

- 1) Create an age variable that can take values from 1 through 8, depending on the age range of the patient. Denote this variable as AGE.
- 2) Create an indicator variable for whether a survey was from the surgical service line. Let this variable equal 1 if surgical and equal to 0 if not surgical. Denote this variable as SURG.
- 3) Create an indicator variable for whether a survey was from the maternity service line. Let this variable equal 1 if maternity and equal to 0 if not maternity. Denote this variable as MAT.
- 4) At this point, every completed survey should have a value from 1 to 8 for AGE, a value of 0 or 1 for SURG, and a value of 0 or 1 for MAT.

The surgical by age interaction variable (Surgical*Age) is equal to the product of SURG and AGE. Similarly, the maternity by age interaction variable (Maternity*Age) is calculated as the product of MAT and AGE. To obtain hospital-level values for these two interaction variables, simply average all the survey-level values just calculated for Surgical*Age and Maternity*Age.

The formula for applying patient mix adjustment is as follows:

If y is the mode-adjusted hospital mean of an HCAHPS outcome (top-box or bottom-box)

a_1 - a_{16} are the individual-level adjustments from Table 1 or Table 2 for the 16 rows other than reference categories (in proportion rather than percentage form)

m_1 - m_{16} are the national means for the PMA variables in the same rows in Table 3

h_1 - h_{16} are the PMA means for the hospital in question in the same form as in Table 3,

then $y' = y + a_1(h_1 - m_1) + a_2(h_2 - m_2) + \dots + a_{16}(h_{16} - m_{16})$ is the patient-mix and mode-adjusted hospital score for that outcome.

For public reporting purposes, HCAHPS scores are rounded to integer percentages. Rounding occurs within top, middle, and bottom-box scores only after patient-mix and mode adjustments have been applied. If the sum of the three scores is not 100%, a further adjustment is made to the middle-box score.

Note: Effective July 1, 2010, the “Point of Origin for Admission or Visit” code for Emergency Room (code 7) was discontinued for use by Medicare Systems. As a result, the patient-mix variable *ER Admission* was not included in the calculation of HCAHPS scores beginning with second quarter 2010, nor will it be included in subsequent quarters. This means the four-quarter average that forms the scores for October 2011 public reporting will contain one quarter that includes ER Admission as a patient-mix adjuster and three quarters that do not. Accordingly, the patient-mix adjustments in Tables 1 and 2 represent the one quarter in which ER Admission was included as an adjuster, and a value of zero for the three quarters in which ER Admission was not included as an adjuster. The national mean for ER Admission reported in Table 3 is based on the quarter for which ER admission was included as an adjuster. The averages and coefficients presented in this document were selected to help hospitals closely approximate their publicly reported HCAHPS scores for the four quarters used in October 2011 public reporting.

Please note: The information presented here will permit a hospital to closely approximate the effect of patient-mix adjustment on its HCAHPS results. However, *exact replication* of published HCAHPS results may not be possible because of (1) the effects of data cleaning and (2) small differences between the effects of quarterly patient-mix adjustments and the 4-quarter averages presented here. For each future public reporting period, Tables 1, 2 and 3 will be updated and will be posted on www.hcahponline.org.

Table 1: “Top-Box” HCAHPS Patient-mix Adjustments
 (Four Quarter Average for October 2011 Public Reporting,
 January 2010 to December 2010 Discharges)

Patient-Mix Adjustment (PMA)	Comm. with Nurses	Comm. with Doctors	Responsiveness of Hosp. Staff	Pain Management	Comm. About Medicines	Cleanliness of Hosp. Env.	Quietness of Hosp. Env.	Discharge Information	Overall Hospital Rating	Recommend the Hospital
Education (per level) (1=8th grade or less and 6=More than 4-year college degree)	2.05%	1.76%	2.61%	2.55%	3.19%	1.97%	3.82%	0.85%	3.25%	1.23%
Self-Rated Health (per level) (1=Excellent and 5=Poor)	5.22%	5.00%	6.39%	6.72%	5.24%	4.09%	4.24%	1.34%	6.36%	5.52%
ER Admission ¹	0.28%	1.09%	0.52%	0.46%	0.65%	0.07%	0.47%	0.77%	0.41%	0.46%
Response Percentile (per 1% of response percentile)	0.21%	0.17%	0.23%	0.16%	0.18%	0.07%	0.04%	0.01%	0.18%	0.17%
Non-English Primary Language	0.31%	-0.49%	0.07%	-1.46%	-1.71%	1.06%	-5.65%	-1.79%	-7.15%	-5.70%
AGE										
Age 18-24	3.64%	3.20%	2.69%	5.21%	-9.95%	4.08%	-8.19%	-2.10%	17.24%	15.41%
Age 25-34	0.78%	0.57%	-3.97%	1.73%	-10.79%	4.23%	-8.19%	-3.48%	13.48%	10.33%
Age 35-44	-1.40%	-1.45%	-6.13%	-0.97%	-12.13%	3.83%	-6.07%	-3.99%	9.34%	7.43%
Age 45-54	-3.49%	-3.00%	-7.34%	-3.23%	-12.38%	3.04%	-3.27%	-4.36%	4.87%	4.13%
Age 55-64	-5.33%	-4.76%	-8.45%	-6.02%	-12.49%	2.24%	-2.75%	-5.02%	0.40%	0.91%
Age 65-74	-5.23%	-5.60%	-7.60%	-6.55%	-10.25%	2.68%	-1.98%	-4.15%	-1.74%	-0.33%
Age 75-84	-3.05%	-3.04%	-4.30%	-3.71%	-5.25%	1.76%	-0.41%	-1.69%	-1.85%	-0.39%
Age 85+ (REFERENCE)	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%
SERVICE LINE										
Maternity	-6.57%	-13.05%	-14.04%	-12.15%	-11.62%	2.33%	-10.34%	-9.00%	-12.37%	-13.89%
Surgical	1.46%	-9.62%	1.47%	-1.72%	-1.32%	0.16%	-0.71%	-6.07%	-4.38%	-3.80%
Medical (REFERENCE)	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%
INTERACTIONS										
Surgical Line * Age ²	-0.09%	0.90%	-0.34%	-0.43%	0.18%	-0.16%	0.04%	-0.10%	0.44%	0.37%
Maternity Line * Age ²	1.08%	1.56%	1.63%	1.69%	2.02%	0.45%	0.74%	1.65%	1.97%	2.28%

¹ Values of zero are averaged in for Q2 2010, Q3 2010, and Q4 2010, so that coefficients are appropriate for approximating the four quarters of October 2011 public reporting

² Age takes on the values of 1 to 8 as follows: (1: 18 to 24); (2: 25 to 34); (3: 35 to 44); (4: 45 to 54); (5: 55 to 64); (6: 65 to 74); (7: 75 to 84); and (8: 85+).

Table 2: “Bottom-Box” HCAHPS Patient-mix Adjustments
 (Four Quarter Average for October 2011 Public Reporting,
 January 2010 to December 2010 Discharges)

Patient-Mix Adjustment (PMA)	Comm. with Nurses	Comm. with Doctors	Responsiveness of Hosp. Staff	Pain Management	Comm. About Medicines	Cleanliness of Hosp. Env.	Quietness of Hosp. Env.	Discharge Information	Overall Hospital Rating	Recommend the Hospital
Education (per level) (1=8th grade or less and 6=More than 4-year college degree)	-0.08%	-0.32%	-0.24%	-0.34%	-1.95%	-0.16%	-1.28%	-0.85%	-0.77%	-0.61%
Self-Rated Health (per level) (1=Excellent and 5=Poor)	-1.92%	-1.89%	-2.83%	-2.52%	-3.72%	-1.79%	-1.84%	-1.34%	-2.81%	-1.77%
ER Admission ¹	-0.15%	-0.46%	-0.28%	-0.25%	-0.52%	-0.07%	-0.29%	-0.77%	-0.25%	-0.16%
Response Percentile (per 1% of response percentile)	-0.05%	-0.05%	-0.08%	-0.04%	-0.11%	-0.01%	0.00%	-0.01%	-0.06%	-0.03%
Non-English Primary Language	0.00%	0.65%	-1.12%	1.09%	3.11%	-0.22%	2.17%	1.79%	2.96%	2.24%
AGE										
Age 18-24	-4.31%	-5.22%	-8.48%	-8.40%	1.70%	-5.98%	-1.92%	2.10%	-8.19%	-7.26%
Age 25-34	-3.33%	-4.18%	-5.68%	-6.87%	2.23%	-5.74%	-1.94%	3.48%	-6.55%	-5.89%
Age 35-44	-2.47%	-3.13%	-4.37%	-5.55%	3.47%	-5.81%	-3.10%	3.99%	-5.36%	-4.77%
Age 45-54	-1.27%	-1.71%	-2.61%	-3.92%	3.86%	-5.35%	-3.67%	4.36%	-3.39%	-3.34%
Age 55-64	0.20%	-0.11%	-0.20%	-1.48%	5.11%	-3.81%	-2.17%	5.02%	-1.06%	-1.61%
Age 65-74	0.80%	0.98%	0.89%	-0.03%	5.05%	-2.83%	-1.10%	4.15%	0.15%	-0.50%
Age 75-84	0.75%	0.85%	1.00%	0.37%	3.31%	-1.30%	-0.71%	1.69%	0.62%	0.15%
Age 85+ (REFERENCE)	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%
SERVICE LINE										
Maternity	4.34%	7.31%	8.82%	7.83%	8.91%	0.67%	6.04%	9.00%	8.18%	6.35%
Surgical	0.32%	5.84%	0.24%	3.11%	1.59%	0.22%	0.42%	6.07%	2.76%	2.16%
Medical (REFERENCE)	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%
INTERACTIONS										
Surgical Line * Age ²	-0.05%	-0.65%	0.12%	-0.18%	-0.10%	0.08%	0.05%	0.10%	-0.35%	-0.32%
Maternity Line * Age ²	-0.56%	-0.85%	-1.06%	-1.03%	-1.40%	-0.41%	-0.26%	-1.65%	-1.09%	-0.94%

¹ Values of zero are averaged in for Q2 2010, Q3 2010, and Q4 2010, so that coefficients are appropriate for approximating the four quarters of October 2011 public reporting

² Age takes on the values of 1 to 8 as follows: (1: 18 to 24); (2: 25 to 34); (3: 35 to 44); (4: 45 to 54); (5: 55 to 64); (6: 65 to 74); (7: 75 to 84); and (8: 85+).

Table 3: National Means of PMA Variables
 (Four Quarter Average for October 2011 Public Reporting,
 January 2010 to December 2010 Discharges)

Patient-Mix Adjustment (PMA)	National Mean
Education (per level) (1=8th grade or less and 6=More than 4-year college degree)	3.655
Self-Rated Health (per level) (1=Excellent and 5=Poor)	2.811
ER Admission ¹	0.450
Response Percentile (per 1% of response percentile)	16.2%
Non-English Primary Language	0.060
AGE	
Age 18-24	0.044
Age 25-34	0.106
Age 35-44	0.074
Age 45-54	0.114
Age 55-64	0.177
Age 65-74	0.213
Age 75-84	0.189
Age 85+ (REFERENCE)	0.082
SERVICE LINE	
Maternity	0.136
Surgical	0.352
Medical (REFERENCE)	0.513
INTERACTIONS	
Surgical Line * Age ²	1.863
Maternity Line * Age ²	0.310

¹ Calculated only from Q110 values, so that coefficients are appropriate for approximating the four quarters of October 2011 public reporting

² Age takes on the values of 1 to 8 as follows: (1: 18 to 24); (2: 25 to 34); (3: 35 to 44); (4: 45 to 54); (5: 55 to 64); (6: 65 to 74); (7: 75 to 84); and (8: 85+).